

2023 PCMH Quality Metrics and Thresholds – Adult Practice

	Core Metric	Definition	Threshold
1.	Antidepressant	The percentage of members 18 years of age and older who were	≥ 40.00%
	Medication Management	treated with antidepressant medication, had a diagnosis of major	
	(adults only) -	depression and who remained on an antidepressant medication	
	Continuation Phase	treatment. Two rates are reported.	
		• Effective Acute Phase Treatment. The percentage of	
		members who remained on an antidepressant	
		medication for at least 84 days (12 weeks).	
		Effective Continuation Phase Treatment. The percentage of	
		members who remained on an antidepressant medication for at	
		least 180 days (6 months).	
2.	Blood Pressure Control	The percentage of members 18–75 years of age with diabetes	≥ 56.00%
	for Patients with Diabetes	(types 1 and 2) whose blood pressure (BP) was adequately	
		controlled (<140/90 mm Hg) during the measurement year.	
3.	Eye Exam for Patients	The percentage of members 18–75 years of age with diabetes	≥ 51.00%
	with Diabetes	(types 1 and 2) who had a retinal eye exam.	
4.	Hemoglobin A1c Control	The percentage of members 18–75 years of age with diabetes	≤ 47.00%
	for Patients with	(types 1 and 2) whose hemoglobin A1c (HbA1c) was at the	
	Diabetes: HbA1c poor	following levels during the measurement year:	
	control (>9.0%)	• HbA1c Control (<8.0%).	
		HbA1c Poor Control (>9.0%).	
5.	Child and Adolescent		
	Well-Care Visits 12 – 21		
	years		
	Child and Adolescent	The percentage of members aged 12-17 years of age who had at	≥ 57.00%
	Well-Care Visits, Ages	least one comprehensive well-care visit with a PCP or an OB/GYN	
	12 – 17 years	practitioner during the measurement year.	
	Child and Adolescent	The percentage of members aged 18-21 years of age who had at	≥ 39.00%
	Well-Care Visits, Ages	least one comprehensive well-care visit with a PCP or an OB/GYN	
	18 – 21 years	practitioner during the measurement year.	